



Acer xfreemanii 'Armstrong'



| | |
|------------------------------------|--|
| Высота | 10 -15 m |
| Ширина | 4-6m |
| Крона | columnar, dark, dense crown |
| Кора и ветви | no particulars |
| Лист | 3/5 lobes, green, 10 -20 cm |
| Осенний окрас | оранжевый, красный |
| Цветы | flower corymbs, red, March/April |
| Плоды | none, fruitless cultivar |
| Колючки/шипы | None |
| Ядовитость | неядовитый, обычно |
| Вид почвы | not too limy |
| Мощение | устойчив к мощению |
| Зона зимостойкости | 4 (-34,4 до -28,9 °C) |
| Устойчивость к ветру | very good |
| Прочие виды устойчивости | устойчив к морозам, зона 1 - 6, сустойчив к ветру |
| Дерево-среда обитания для животных | устойчив к морозам, зона 1 - 6, сустойчив к ветру, притягивает бабочек |
| Применение | аллеи и широкие улицы, узкие улицы |
| Форма | штамбовое дерево, многоствольное дерево |
| Происхождение | Scanlon Nurseries, Ohio, USA, 1951 |
| Синонимы | Acer rubrum 'Armstrong' |

This cultivar originated in the USA in 1951 and until recently was considered to belong to A. rubrum. Recent studies reveal that 'Armstrong' came into being from a crossing between A. rubrum and A. saccharinum and therefore belongs to the hybrid Acer x freemanii. This hybrid was not only created by man, for it also grows in the wild in the USA. The green leaves have 3 to 5 lobes. Their under surface is grey-green. In our country the beautiful orange-red autumn colour can only be seen after a hot summer. It roots widely with a dense branching. 'Armstrong' requires a not too limy soil. The tree can be used well in hard surfaces, which renders it very suitable to be used in narrow streets and avenues.