

https://www.vdberk.ru/index.php/derevya/quercus-coccinea-splendens/



Quercus coccinea 'Splendens'



| Высота | 8 m |
|--------------------------|--|
| Ширина | 12-15m |
| Крона | flat-topped spherical, half-open crown |
| Кора и ветви | twigs brown, with thorns, bark grey-brown, slightly grooved |
| Лист | broad oval, glossy dark green, 5 - 10 cm, autumn colour yellow, orange and red |
| Осенний окрас | красный |
| Цветы | white, in felt-like panicles, May-June |
| Плоды | round, bright red, Ø about 1 cm |
| Колючки/шипы | None |
| Ядовитость | неядовитый, обычно |
| Вид почвы | cannot tolerate wet soil |
| Влажность почвы | устойчив к влажным почвам |
| Мощение | неустойчив к мощению |
| Зона зимостойкости | 5b (-26,0 до -23,4 °C) |
| Устойчивость к ветру | good |
| Прочие виды устойчивости | устойчив к морозам, зона 1 - 6, сустойчив к ветру, устойчив к соли для улиц |
| Применение | аллеи и широкие улицы, промышленные зоны |
| Форма | штамбовое дерево, многоствольное дерево |
| Происхождение | Knap Hill Nursery, Woking, England, circa 1900 |
| | |

Forms a broad crown with branches growing horizontally. The bark is dark grey to almost black and is moderately rough. Q. coccinea bears a strong resemblance to Q. palustris. But Q. coccinea's young twigs are yellowish brown while those of Q. palustris are olive-green. The leaves are elongated to oval (sometimes obovate) and pinnated with 7 - 9 projecting serrated lobes. They are less deeply incised than those of Q. palustris and are 9 -15 cm long and 6 - 11 (15) cm wide. In the autumn the leaves turn a deep scarlet red. This Quercus coccinea selection is distinguished from the species by its even more intense autumnal colour. Once the leaves have turned brown they remain on the tree until halfway through winter. The acorns are single and one-third enclosed by the cupule.