

https://www.vdberk.ru/index.php/derevya/ulmus-minor/



Ulmus minor



Высота	20 - 30 m, fast growing
Ширина	12-15m
Крона	broad fan-shaped to broad oval, half-open crown, capricious growing
Кора и ветви	has brownish-grey grooves, flakes off in platelets, twigs brown
Лист	oval-elongated to obovate, glossy green, 5 - 10 cm
Цветы	in ver dense bundles, purplish-red, March, before the leaves emerge
Плоды	winged nutlet, 1.5 - 2 cm
Колючки/шипы	None
Ядовитость	неядовитый, обычно
Вид почвы	moist and light, favours calcareous soils
Влажность почвы	устойчив к влажным почвам, устойчив к коротким наводнениям, устойчив к длительным наводнениям
Мощение	неустойчив к мощению
Зона зимостойкости	5 (-28,8 до -23,4 °C)
Устойчивость к ветру	very good, also tolerates sea wind
Прочие виды устойчивости	устойчив к морозам, зона 1 - 6, сустойчив к ветру
Дерево-среда обитания для животных	устойчив к морозам, зона 1 - 6, сустойчив к ветру, притягивает бабочек
Применение	аллеи и широкие улицы, побережье, промышленные зоны
Форма	штамбовое дерево, многоствольное дерево
Происхождение	western Europe, north Africa, south-west Asia
Синонимы	Ulmus carpinifolia, Ulmus campestris

Fast-growing tall tree with a dense crown and a variable manner of growth, from broad fan-shaped to broad oval. Occasionally a cork layer occurs on the bare twigs. The leaves are oval-elongated to obovate, widest in or above the centre. On the upper side they are smooth and glossy green, on the underside practically bare. The foot of the leaf covers one side of the short stalk, a characteristic of elm leaves. The leaf edge is double-serrated. The flowers grow in very dense bundles of 15 to 30 together. They appear before the leaves emerge and are purplish-red. The winged fruits are also often present before the emergence of the leaves: the fruits are then still green but later turn brown. It is fairly badly sensitive to Dutch elm disease. Ulmus minor used to be planted a great deal but is very little cultivated at present.