



Acer platanoides 'Reitenbachii'



Высота	15-20 m
Ширина	10-12m
Крона	broad ovoid, dark, dense crown
Кора и ветви	grey
Лист	green, redbrown
Цветы	yellowgreen, flowers in April
Плоды	brown, inconspicuous
Колючки/шипы	None
Ядовитость	ядовитые части
Вид почвы	clayed soil, loamy soil, sandy soil
Влажность почвы	устойчив к коротким наводнениям
Мощение	устойчив к частичному мощению
Зона зимостойкости	4 (-34,4 до -28,9 °C)
Устойчивость к ветру	not very sensitive, not resistant to sea wind
Прочие виды устойчивости	устойчив к морозам, зона 1 - 6
Дерево-среда обитания для животных	устойчив к морозам, зона 1 - 6, притягивает бабочек
Применение	аллеи и широкие улицы
Форма	штамбовое дерево

This tree's history goes back to the nursery of J. Reitenbach in Poland, at that time part of Russia. The plant was named by Johann Xaver Robert Caspary, a German botanist, but it was introduced to the market by the Belgian nursery of Louis Van Houtte. In terms of habit, 'Reitenbachii' is typical of the species. The sprouting leaf is a shiny dark brown, turning to matt dark green with dark red veins in summer. An attractive tree, somewhat resembling 'Schwedleri', which has more intense leaf colours. Grows on any soil, but does not tolerate stagnating groundwater and peaty soils.