

https://www.vdberk.ru/derevya/koelreuteria-paniculata-september/



## Koelreuteria paniculata 'September'



Высота	7 - 17 m
Ширина	7-15m
Крона	broad, almost round, somewhat irregular, light, open crown
Кора и ветви	grey-brown, slightly grooved
Лист	(doubly) pinnate, dull green, 30 - 35 cm, feathered leaf
Осенний окрас	желтый, оранжевый
Цветы	yellow, in large terminal plumes, August/September
Плоды	bladder-shaped, bronze in colour, seeds black, Ø approx. 4.5 cm
Колючки/шипы	None
Ядовитость	неядовитый, обычно
Вид почвы	makes very few demands on the soil, stands up to dry conditions
Влажность почвы	устойчив к сухим почвам
Мощение	неустойчив к мощению
Зона зимостойкости	6b (-20,5 до -17,8 °C)
Устойчивость к ветру	moderate
Прочие виды устойчивости	устойчив к морозам, зона 1 - 6
Дерево-среда обитания для животных	устойчив к морозам, зона 1 - 6, притягивает бабочек
Применение	кашпо для деревьев
Форма	штамбовое дерево, многоствольное дерево
Происхождение	Indiana University, Bloomington, USA, 1960

This American selection initially shows a slower growth rate than the species. The rounded crown grows steadily broader. Widely-spread branches bend downwards. The large, pinnate (sometimes doubly pinnate) leaves consist of 7 - 15 leaf segments. These are lobed, approx. 7 - 10 cm in length and have a serrated edge. The leaves are a bronze colour in the spring and turn yellow to orange in the autumn. The numerous small, yellow flowers are in terminal plumes approx. 30 cm long. The tree flowers 6 - 8 weeks later than the species and the flowers are often very numerous. The bladder-like fruit, approx 4.5 cm long, change colour from green to bronze and contain a few black seeds. The fruit remains on the tree for a long time. This tree prefers a sheltered site since young specimens can suffer frost damage, but recover well.