



Quercus robur 'Timuki'



Высота	15 - 20 m
Ширина	8-12m
Крона	round, half-open crown
Кора и ветви	twigs dark brown, bark dark grey, deeply grooved
Лист	obovate, lobed, dark greenish purple, 7 - 14 cm
Цветы	green-yellow catkins, 2 - 4 cm, May
Плоды	ovate to oval, 1.5 - 2.5 cm, in groups of 2 - 6, cupule with triangular contiguous scales
Колючки/шипы	None
Ядовитость	неядовитый, обычно
Вид почвы	fertile and moist
Влажность почвы	устойчив к влажным почвам
Мощение	неустойчив к мощению
Зона зимостойкости	5a (-28,8 до -26,1 °C)
Устойчивость к ветру	good
Прочие виды устойчивости	устойчив к морозам, зона 1 - 6, устойчив к ветру, устойчив к соли для улиц
Применение	промышленные зоны
Форма	штамбовое дерево
Происхождение	Raimond Cinovskis, Riga, Latvia, 1946

Red-leaved cultivar of *Q. robur* with a broad ovoid crown, eventually becoming round. Grows significantly better than the old purple-leaved cultivar, 'Purpurascens'. The bark, blossom and fruit are the same as the species. The obovate leaves are, in terms of shape, also like the species although they are larger than on 'Purpurascens'. On emerging the leaf is green but it gradually turns darker, and at the end of summer it turns through red to a glossy dark greenish purple. The tree was found by M. Timuki in 1939 but was only named and distributed in 1946. The good growth and lovely foliage colours are the most important features of this purple-leaved oak. In addition, 'Timuki' is significantly less susceptible to mildew.